

measurement ERROR webinar series

Recommended Resources

- Bailey RL, Dodd KW, Goldman JA, Gahche JJ, Dwyer JT, Moshfegh AJ, Sempos CT, Picciano MF. Estimation of total usual calcium and vitamin D intakes in the United States. *J Nutr.* 2010;140(4):817-22.
- Carroll RJ, Midthune D, Subar AF, Shumakovich M, Freedman LS, Thompson FE, Kipnis V. Taking advantage of the strengths of two different dietary assessment instruments to improve intake estimates for nutritional epidemiology. *Am J Epidemiol.* In press.
- Carroll RJ, Ruppert D, Stefanski LA, Crainiceanu CM. *Measurement Error in Nonlinear Models: A Modern Perspective*, 2nd ed. Boca Raton, FL: Chapman & Hall/CRC, 2006.
- Dodd KW, Guenther PM, Freedman LS, Subar AF, Kipnis V, Midthune D, Tooze JA, Krebs-Smith SM. Statistical methods for estimating usual intake of nutrients and foods: a review of the theory. *J Am Diet Assoc.* 2006;106(10):1640-50.
- Freedman LS, Kipnis V, Schatzkin A, Tasevska N, Potischman N. Can we use biomarkers in combination with self-reports to strengthen the analysis of nutritional epidemiologic studies? *Epidemiol Perspect Innov.* 2010;7(1):2.
- Freedman LS, Midthune D, Carroll RJ, Tasevska N, Schatzkin A, Mares J, Tinker L, Potischman N, Kipnis V. Using regression calibration equations that combine self-reported intake and biomarker measures to obtain unbiased estimates and more powerful tests of dietary associations. *Am J Epidemiol.* In press.
- Freedman LS, Schatzkin A, Midthune D, Kipnis V. Dealing with dietary measurement error in nutritional cohort studies. *J Natl Cancer Inst.* 2011;103(14):1086-92.
- Freedman LS, Tasevska N, Kipnis V, Schatzkin A, Mares J, Tinker L, Potischman N. Gains in statistical power from using a dietary biomarker in combination with self-reported intake to strengthen the analysis of a diet-disease association: an example from CAREDS. *Am J Epidemiol.* 2010;172(7):836-42.
- Garriguet D. Combining nutrient intake from food/beverages and vitamin/mineral supplements. *Health Rep.* 2010;21(4):71-84.
- Guenther PM, Reedy J, Krebs-Smith SM, Reeve BB. Evaluation of the Healthy Eating Index-2005. *J Am Diet Assoc.* 2008;108(11):1854-64.
- Guenther PM, Reedy J, Krebs-Smith SM. Development of the Healthy Eating Index-2005. *J Am Diet Assoc.* 2008;108(11):1896-901.
- Institute of Medicine. *Dietary Reference Intakes: Applications in Dietary Assessment.* Washington, DC: National Academy Press, 2000.
- Kipnis V, Freedman LS, Brown CC, Hartman A, Schatzkin A, Wacholder S. Interpretation of energy adjustment models for nutritional epidemiology. *Am J Epidemiol.* 1993;137:1376-80.
- Kipnis V, Midthune D, Buckman DW, Dodd KW, Guenther PM, Krebs-Smith SM, Subar AF, Tooze JA, Carroll RJ, Freedman LS. Modeling data with excess zeros and measurement error: application to evaluating relationships between episodically consumed foods and health outcomes. *Biometrics.* 2009;65(4):1003-10.
- Kipnis V, Subar AF, Midthune D, Freedman LS, Ballard-Barbash R, Troiano RP, Bingham S, Schoeller DA, Schatzkin A, Carroll RJ. Structure of dietary measurement error: results of the OPEN biomarker study. *Am J Epidemiol.* 2003;158(1):14-21; discussion 22-6.
- Korn EL, Graubard BI. *Analysis of Health Surveys.* New York, NY: John Wiley & Sons; 1999.

- Midthune D, Schatzkin A, Subar AF, Thompson FE, Freedman LS, Carroll RJ, Shumakovich MA, Kipnis V. Validating an FFQ for intake of episodically consumed foods: application to the National Institutes of Health-AARP Diet and Health Study. *Public Health Nutr.* 2011; 13:1-10.
- National Cancer Institute usual intakes web pages: <http://riskfactor.cancer.gov/diet/usualintakes/>
- National Health and Nutrition Examination Survey (NHANES) Dietary Tutorial: <http://www.cdc.gov/nchs/tutorials/Dietary/index.htm>.
- National Research Council. *Nutrient Adequacy. Assessment Using Food Consumption Surveys.* Washington, DC: National Academy Press; 1986.
- Park SY, Murphy SP, Wilkens LR, Yamamoto JF, Kolonel LN. Allowing for variations in multivitamin supplement composition improves nutrient intake estimates for epidemiologic studies. *J Nutr.* 2006;136(5):1359-64.
- Reedy J, Mitrou PN, Krebs-Smith SM, Wirfält E, Flood A, Kipnis V, Leitzmann M, Mouw T, Hollenbeck A, Schatzkin A, Subar AF. Index-based dietary patterns and risk of colorectal cancer: the NIH-AARP Diet and Health Study. *Am J Epidemiol.* 2008;168(1):38-48.
- Rosner B, Spiegelman D, Willett WC. Correction of logistic regression relative risk estimates and confidence intervals for measurement error: the case of multiple covariates measured with error. *Am J Epidemiol.* 1990;132(4):734-45.
- Rosner B, Willett WC, Spiegelman D. Correction of logistic regression relative risk estimates and confidence intervals for systematic within-person measurement error. *Stat Med.* 1989;8(9):1051-69; discussion 1071-3.
- Schatzkin A, Subar AF, Moore S, Park Y, Potischman N, Thompson FE, Leitzmann M, Hollenbeck A, Morrissey KG, Kipnis V. Observational epidemiologic studies of nutrition and cancer: the next generation (with better observation). *Cancer Epidemiol Biomarkers Prev.* 2009;18(4):1026-32.
- Thiébaud AC, Kipnis V, Schatzkin A, Freedman LS. The role of dietary measurement error in investigating the hypothesized link between dietary fat intake and breast cancer – a story with twists and turns. *Cancer Invest.* 2008;26(1):68-73.
- Thompson FE, Subar AF. Dietary assessment methodology, In: Coulston AM, Boushey CJ, editors, *Nutrition in the prevention and treatment of disease*, 2nd ed. San Diego, CA: Academic Press. 2008:3-39. (Also available online at: (http://riskfactor.cancer.gov/diet/adi/thompson_subar_dietary_assessment_methodology.pdf))
- Toozé JA, Kipnis V, Buckman DW, Carroll RJ, Freedman LS, Guenther PM, Krebs-Smith SM, Subar AF, Dodd KW. A mixed-effects model approach for estimating the distribution of usual intake of nutrients: the NCI method. *Stat Med.* 2010;29(27):2857-68.
- Toozé JA, Midthune D, Dodd KW, Freedman LS, Krebs-Smith SM, Subar AF, Guenther PM, Carroll RJ, Kipnis V. A new statistical method for estimating the usual intake of episodically consumed foods with application to their distribution. *J Am Diet Assoc.* 2006;106(10):1575-87.
- Willett WC, Howe GR, Kushi LH. Adjustment for total energy intake in epidemiologic studies. *Am J Clin Nutr.* 1997;65(4 Suppl):1220S-1228S; discussion 1229S-1231S.
- Zhang S, Krebs-Smith SM, Midthune D, Perez A, Buckman D, Kipnis V, Freedman LS, Dodd KW, Carroll RJ. Fitting a bivariate measurement error model for episodically consumed dietary components. *Intl J Biostat.* 2011;7(1):Article 1.
- Zhang S, Midthune D, Guenther PM, Krebs-Smith SM, Kipnis V, Dodd KW, Buckman DW, Toozé JA, Freedman L, Carroll RJ. A new multivariate measurement error model with zero-inflated dietary data, and its application to dietary assessment. *Ann Appl Stat.* 2011;5(2B):1456-87.